

Polynomial equations (exercises from class and more)

Mr. Neeman. 10A, August 11, 2011

Here are the equations we've seen so far in class, plus some others. Afterwards are the answers and in some cases comments about how they can be solved.

#1. $x^3 + x^2 - 2x - 2 = 0$

#2. $x^4 - x^3 + 4x^2 - 4x = 0$

#3. $x^3 - x^2 - 10x - 8 = 0$

#4. $x^3 + 3x^2 + x - 2 = 0$

#5. $x^3 + 3x^2 + x - 2 = 0$

#6. $x^3 - 6x + 3x + 10 = 0$

#7. $2x^3 + 10x^2 + 2x - 30 = 0$

#8. $x^4 - x^2 - 12 = 0$

#9. $x^4 + 2x^3 - 2x^2 - 4x - 16 = 0$

#10. $x^3 + 4x^2 + 10x + 12 = 0$

#11. $x^4 + 4x^3 + 7x^2 - 4x - 8 = 0$

#12. $x^3 - 4x^2 - 29x + 12 = 0$

#13. $3x^4 + 3x^3 + 6x^2 + 6x = 0$

#14. $x^3 + 4x^2 - 11x - 30 = 0$

Answers

#1. $x^3 + x^2 - 2x - 2 = 0$

You can find the root $x = -1$ by trying the divisors of 2, and after factoring it out are left with $x^2 - 2$ which you can factorize.

$(x + 1)(x - \sqrt{2})(x + \sqrt{2}) = 0$. So $x \in \{-1, \sqrt{2}, -\sqrt{2}\}$

#2. $x^4 - x^3 + 4x^2 - 4x = 0$

You can first factor out an x , which means that $x = 0$ is a root. You can then test the divisors of 4 and see $x = 1$ works. So you can factor out $x - 1$ which leaves $x^2 + 4$. Since this quadratic has no solutions, the two solutions above are the only ones.

$x(x^2 + 4)(x - 1) = 0$. So $x \in \{0, 1\}$

#3. $x^3 - x^2 - 10x - 8 = 0$

$(x + 2)(x + 1)(x - 4) = 0$. So $x \in \{-2, -1, 4\}$

#4. $x^3 + 3x^2 + x - 2 = 0$

You can find $x = -4$, and after you factor out $x + 4$ you get $x^2 + 1$ which has no solutions.

$(x + 4)(x^2 + 1) = 0$. So $x = -4$

#5. $x^3 + 3x^2 + x - 2 = 0$

You can find $x = -2$ and after factoring out $x + 2$ you get $x^2 + x - 1$ left, which has irrational roots $\frac{1 \pm \sqrt{5}}{2}$. So $x \in \{-2, \frac{-1 - \sqrt{5}}{2}, \frac{-1 + \sqrt{5}}{2}\}$

#6. $x^3 - 6x + 3x + 10 = 0$

$(x - 5)(x + 1)(x - 2) = 0$. So $x \in \{-1, 2, 5\}$

#7. $2x^3 + 10x^2 + 2x - 30 = 0$

First, you can divide both sides of the equation by 2. Then you can try the divisors of 15, finding that $x = -3$ is a solution. You can then factor out $x + 3$ and have left $x^2 + 2x - 5$ which has two irrational solutions.

$2(x + 3)(x + 1 - \sqrt{6})(x + 1 + \sqrt{6}) = 0$. So $x \in \{-3, -1 - \sqrt{6}, -1 + \sqrt{6}\}$

#8. $x^4 - x^2 - 12 = 0$

This one can be done by the usual method, trying out the divisors of 12. You find $x = 2$ and $x = -2$ are roots, and after factoring out by $x - 2$ and $x + 2$ you're left with a quadratic with no solutions. However, one could also take the equation as a quadratic in x^2 : $(x^2)^2 - x^2 - 12 = 0$, and factorize this by inspection, getting $(x^2 - 4)(x^2 + 3) = 0$, from which you can then find the solutions.

$(x + 2)(x - 2)(x^2 + 3) = 0$. So $x \in \{-2, 2\}$.

#9. $x^4 + 2x^3 - 2x^2 - 4x - 16 = 0$

You can test the divisors of 16 to find $x = 2$ and $x = -2$ are solutions. After factoring out $x - 2$ and $x + 2$ you're left with $x^2 + 2x + 4$ which has no solutions.

$(x^2 + 2x + 4)(x + 2)(x - 2) = 0$. So $x \in \{-2, 2\}$.

#10. $x^3 + 4x^2 + 10x + 12 = 0$

You can test the divisors to find $x = -2$ and are left with a quadratic with no solutions.

$(x + 2)(x^2 + 2x + 6) = 0$. So $x = -2$.

#11. $x^4 + 4x^3 + 7x^2 - 4x - 8 = 0$

You can test the divisors of 8 and find $x = -1$ and $x = 1$ are solutions. If you factor out the corresponding linear terms you get a quadratic without solutions.

$(x + 1)(x - 1)(x^2 + 4x + 8) = 0$. So $x \in \{-1, 1\}$.

#12. $x^3 - 4x^2 - 29x + 12 = 0$

You can test the divisors to find $x = -4$ is a solution and after factoring $x + 4$ out you get a quadratic with irrational roots.

$(x + 4)(x - 4 - \sqrt{13})(x - 4 + \sqrt{13}) = 0$. So $x \in \{-4, 4 - \sqrt{13}, 4 + \sqrt{13}\}$.

#13. $3x^4 + 3x^3 + 6x^2 + 6x = 0$

Here you can begin by factoring out $3x$. This tells you either $x = 0$ or $x^3 + x^2 + 2x + 2 = 0$. For this cubic equation, you can try the divisors of 2 and find $x = -1$ is a solution. If you then divide by $x + 1$ you get $x^2 + 2$ which has no solutions. You could also factorize the cubic by grouping: $x^3 + x^2 + 2x + 2 = x^2(x + 1) + 2(x + 1) = (x^2 + 2)(x + 1)$.

$3x(x + 1)(x^2 + 2) = 0$. So $x \in \{-1, 0\}$.

#14. $x^3 + 4x^2 - 11x - 30 = 0$

You can try the divisors of 30. Since there are a lot of them it's a good idea to start with the small ones and when you find one factor out the corresponding linear term.

$(x + 2)(x - 3)(x + 5) = 0$. So $x \in \{-5, -2, 3\}$.